

Gonadotropin-Releasing Hormone Agonists During Chemotherapy for Preservation of Ovarian Function and Fertility in Premenopausal Patients With Early Breast Cancer: A Systematic Review and Meta-Analysis of Individual Patient–Level Data

Matteo Lambertini, Halle C.F. Moore, Robert C.F. Leonard, Sibylle Loibl, Pamela Munster, Marco Bruzzone, Luca Boni, Joseph M. Unger, Richard A. Anderson, Keyur Mehta, Susan Minton, Francesca Poggio, Kathy S. Albain, Douglas J.A. Adamson, Bernd Gerber, Amy Cripps, Gianfilippo Bertelli, Sabine Seiler, Marcello Ceppi, Ann H. Partridge, and Lucia Del Mastro

Author affiliations and support information (if applicable) appear at the end of this article.

Published at jco.org on May 2, 2018.



Processed as a Rapid Communication manuscript.

A.H.P. and L.D.M. are co-last authors.

Clinical trial information: PROSPERO registration number CRD42014015638.

Corresponding author: Matteo Lambertini, MD, Institut Jules Bordet and Université Libre de Bruxelles, Boulevard de Waterloo 121, 1000 Brussels, Belgium; e-mail: matteo.lambertini85@gmail.com.

© 2018 by American Society of Clinical Oncology

0732-183X/18/3699-1/\$20.00

A B S T R A C T

Purpose

The role of temporary ovarian suppression with gonadotropin-releasing hormone agonists (GnRHa) during chemotherapy as a strategy to preserve ovarian function and fertility in premenopausal women remains controversial. This systematic review and meta-analysis using individual patient–level data was conducted to better assess the efficacy and safety of this strategy in patients with early breast cancer.

Methods

The trials in which premenopausal women with early breast cancer were randomly assigned to receive (neo)adjuvant chemotherapy alone or with concurrent GnRHa were eligible for inclusion. Primary end points were premature ovarian insufficiency (POI) rate and post-treatment pregnancy rate. Disease-free survival and overall survival were secondary end points. Because each study represents a cluster, statistical analyses were performed using a random effects model.

Results

A total of 873 patients from five trials were included. POI rate was 14.1% in the GnRHa group and 30.9% in the control group (adjusted odds ratio, 0.38; 95% CI, 0.26 to 0.57; $P < .001$). A total of 37 (10.3%) patients had at least one post-treatment pregnancy in the GnRHa group and 20 (5.5%) in the control group (incidence rate ratio, 1.83; 95% CI, 1.06 to 3.15; $P = .030$). No significant differences in disease-free survival (adjusted hazard ratio, 1.01; 95% CI, 0.72 to 1.42; $P = .999$) and overall survival (adjusted hazard ratio, 0.67; 95% CI, 0.42 to 1.06; $P = .083$) were observed between groups.

Conclusion

Our findings provide evidence for the efficacy and safety of temporary ovarian suppression with GnRHa during chemotherapy as an available option to reduce the likelihood of chemotherapy-induced POI and potentially improve future fertility in premenopausal patients with early breast cancer.

J Clin Oncol 36. © 2018 by American Society of Clinical Oncology

INTRODUCTION

Breast cancer is the most commonly diagnosed malignancy in premenopausal women, and its treatment often results in long-term sequelae and impaired quality of life.¹ Given the improved prognosis of patients with breast cancer over the past years, survivorship issues are becoming more important.¹ The use of anticancer therapies in premenopausal patients with breast cancer is

associated with gonadotoxicity.² Age of the patient at the time of treatment, type of chemotherapy regimen administered, and use of adjuvant endocrine therapy are crucial factors affecting the risk of developing this side effect.² Chemotherapy-induced premature ovarian insufficiency (POI) can have a substantial negative impact on patients' quality of life and is associated with several side effects, such as vasomotor symptoms, sexual dysfunction, and fertility-related problems.³

ASSOCIATED CONTENT



See accompanying Editorial
DOI: <https://doi.org/10.1200/JCO.2018.78.9347>



Data Supplement
DOI: <https://doi.org/10.1200/JCO.2018.78.0858>

DOI: <https://doi.org/10.1200/JCO.2018.78.0858>

International guidelines recommend to counsel all young patients newly diagnosed with breast cancer about the potential risk of chemotherapy-induced POI and infertility.^{4,5} Failure to address these issues can negatively influence patients' psychosocial health and their adherence to the proposed anticancer treatments, potentially affecting disease-related morbidity and mortality.¹ Embryo and oocyte cryopreservation are the current standard strategies for fertility preservation in young women with breast cancer.^{4,5} However, these approaches do not prevent the risk of developing chemotherapy-induced POI. To date, temporary ovarian suppression obtained by administering gonadotropin-releasing hormone agonists (GnRHa) during chemotherapy is the only medical intervention with the potential to preserve ovarian function in premenopausal patients receiving cytotoxic systemic therapy.² Because of the conflicting results reported in randomized studies, the role of this option remains controversial, and it is still considered an experimental technique by major international guidelines.^{4,5}

In 2015, we performed a systematic review and meta-analysis on the basis of abstracted data from publications to investigate the protective role of temporary ovarian suppression with GnRHa during chemotherapy in premenopausal patients with early breast cancer.⁶ The use of GnRHa was associated with a significantly reduced risk of chemotherapy-induced POI and amenorrhea 1 year after chemotherapy completion, as well as an increased chance of obtaining a subsequent pregnancy.⁶ Nevertheless, no final conclusions could be drawn, mainly because of the lack of data on ovarian function beyond 1 year after the end of chemotherapy, the limited information on the safety of this approach, and the lack of availability at that time of the results of the Anglo Celtic Group OPTION (Ovarian Protection Trial in Oestrogen Non-responsive Premenopausal Breast Cancer Patients Receiving Adjuvant or Neo-adjuvant Chemotherapy) trial,⁷ one of the largest studies in this field. Furthermore, without individual patient-level data, analyses of other efficacy and safety outcomes as well as the evaluation of the association between treatment effect and patient or tumor characteristics were not possible. The current study seeks to provide more conclusive clinical evidence on this controversial topic by conducting a meta-analysis on the basis of individual patient-level data of the randomized trials that investigated the role of temporary ovarian suppression with GnRHa during chemotherapy as a strategy to preserve ovarian function and fertility in premenopausal patients with early breast cancer.

METHODS

This systematic review and meta-analysis of individual patient-level data are reported according to the Preferred Reporting Items for Systematic Reviews and Meta-analysis guidelines.⁸ A protocol was developed before study initiation and submitted to PROSPERO (registration number CRD42014015638).

Identification of Studies and Collection of Data

Details about the systematic review of the literature were previously reported.⁶ Briefly, a search using PubMed, Embase, and the Cochrane Library was conducted without any date or language restrictions up to April 30, 2015. Furthermore, conference proceedings presented at the most important international conferences from 2004 onward until April 2015 were searched to identify unpublished studies; cross-referencing from relevant studies and review articles was also conducted to confirm retrieval

of all possible pertinent trials. The search strategy was then repeated before final analysis on August 31, 2017 to confirm the retrieval of all possible trials. Eligible studies were randomized trials evaluating the efficacy of adding GnRHa to chemotherapy as a strategy to reduce the occurrence of chemotherapy-induced POI in premenopausal women with early breast cancer receiving (neo)adjuvant chemotherapy.

For all participants enrolled in each of the included trials, individual patient-level data (baseline patient and tumor characteristics, administered treatments, and data on ovarian function after chemotherapy, pregnancies after breast cancer diagnosis, adverse events during treatment, and survival outcomes) were collected. Data from each of the included trials were carefully checked and verified for consistency with their original publications; discrepancies were discussed and resolved with the authors before pooling the data in the final unified database used for analysis.

Outcomes

This study aimed to evaluate both the efficacy (ie, preservation of ovarian function and fertility) and the safety (ie, toxicity and survival outcomes) of temporary ovarian suppression with GnRHa during chemotherapy in premenopausal patients with early breast cancer. Primary end points were POI rate (according to the definition used as primary end point in each trial) and post-treatment pregnancy rate. Secondary end points included amenorrhea rates 1 year and 2 years after the end of chemotherapy, GnRHa-related adverse events (hot flashes, sweating, mood changes, vaginal dryness, and headache), disease-free survival (DFS), and overall survival (OS). Prespecified subgroup analyses investigated the efficacy and safety of temporary ovarian suppression with GnRHa during chemotherapy according to age of the patients, estrogen receptor status, type and duration of chemotherapy administered, and tumor stage.

Statistical Analysis

All analyses were performed including the total number of patients with available information for each specific end point.

For POI rate, the primary end point definition of POI used in each of the included trials was used. To apply a more homogeneous definition, amenorrhea rates (defined as absence of menses) at 1 year and 2 years after the end of chemotherapy were also computed. For post-treatment pregnancy rate, only the first reported pregnancy for each patient was considered independently of its outcome. Adjusted odds ratios (ORs) with 95% CIs were calculated to estimate the effect size of temporary ovarian suppression with GnRHa during chemotherapy. Fisher's exact test was applied for POI and amenorrhea analyses. Incidence rate ratio (IRR) between the GnRHa and control groups for post-treatment pregnancy rate was computed. A multivariate logistic regression analysis was performed to investigate the effects of GnRHa treatment on the risk of developing POI, 1-year and 2-year amenorrhea adjusting by age of the patients at the time of study entry, estrogen receptor status, type, and duration of chemotherapy administered. Because each trial represents a cluster of allegedly correlated outcomes, a generalized linear mixed model for binary end points with logit link was fitted to the data by adding to the model the random effect of the study. This model allowed estimating the amount of heterogeneity between trials and, accordingly, it provided suitable estimates of the standard errors of the predictors (fixed effects) included in the model.

GnRHa-related adverse events were dichotomized first as no adverse event (grade 0) and adverse event of any grade (1 to 4), and then as no adverse event (grade 0), mild (grade 1 or 2), and severe (grade 3 or 4) adverse events. Toxicity rates were compared using Fisher's exact test.

The reverse Kaplan-Meier method was used to calculate the median period of follow-up and its interquartile range (IQR). DFS interval was computed as the difference between the date of random assignment and the date of locoregional, contralateral, or distant recurrence, second malignancy, or death, whichever occurred first. OS was defined as the time interval between the date of random assignment and the date of death from any cause. Observation times of patients without the event were censored on the date of their last follow-up visit. DFS and OS probabilities were

computed according to the Kaplan-Meier method. To investigate the effect of GnRH_a treatment on the risk of developing DFS and OS events adjusting by age of the patients at the time of study entry, estrogen receptor status, type and duration of chemotherapy administered, and tumor stage, a mixed effect Cox proportional hazards model was fitted to the data to take into account the clustering effect of each study. As estimates of treatment effect, adjusted hazard ratios (HRs) with 95% CI were computed. To check for the proportional hazards assumption, the Schoenfeld residuals were examined.

Subgroup analyses of both efficacy and safety end points were performed by means of an interaction test to determine the consistency of the treatment effect on the outcomes according to age of the patients, estrogen receptor status, type and duration of chemotherapy administered, and tumor stage. Likelihood ratio test was applied to test both the main effects and the interaction effects of the covariates included in the statistical models. In addition, a meta-analysis procedure was applied to the data of each individual study for every end point to evaluate the consistency of the results between trials. As overall measure of the effect across studies, we computed the weighted mean of the OR_{*i*} or HR_{*i*} estimated from each *i*-th trial, with weights proportional to the variance of OR_{*i*} or HR_{*i*}. The meta-analysis was performed by means of the inverse-variance method transforming the OR_{*i*} or HR_{*i*} in its natural logarithm. The heterogeneity between studies was quantified through the Higgins I² index. In the presence of significant heterogeneity, the random effect model following the method

of DerSimonian and Laird was applied, because it is generally more appropriate in such situation.

All statistical tests were two-sided, and *P* values < .05 were considered statistically significant. Statistical analyses were performed using STATA 14.2 (StataCorp LP, College Station, TX).

RESULTS

Of the 676 entries returned by the initial database search, 662 were excluded because they did not meet the inclusion criteria (Data Supplement). A total of 14 publications corresponding to 13 different randomized trials were considered eligible for this study.^{7,9-21} Individual patient-level data were available for five major trials (PROMISE-GIM6 [PRevention Of Menopause Induced by chemotherapy: A Study in Early breast cancer patients—Gruppo Italiano Mammella 6],^{9,10} POEMS [Prevention Of Early Menopause Study]/SWOG S0230,¹¹ Anglo Celtic Group OPTION,⁷ GBG [German Breast Group]-37 ZORO [ZOladex Rescue of Ovarian function],¹² Moffitt-led trial¹³) including 873 randomly assigned patients. Individual patient-level data from 708 patients included in the remaining

Table 1. Characteristics of the Five Included Randomized Trials

Characteristic	PROMISE-GIM6 ^{9,10}	POEMS/SWOG S0230 ¹¹	Anglo Celtic Group OPTION ⁷	GBG-37 ZORO ¹²	Moffitt-led trial ¹³
Type of chemotherapy	CMF or E→CMF or EP→CMF or ED→CMF or AC or EC or FEC or AC→D or EC→D or EC→P or FEC→P or FEC→D or ED	AC or CAF or TAC or CEF or AC→T or CMF	CAF or CAF→T or CEF or CEF→T	FEC→D or EC→D or FEC or FAC or TAC or EC→D or AC→D or FEC→GEM	AC or AC→T or FEC or FAC
No. of cycles of chemotherapy	6-8	3-8	6-8	6-8	4-8
Definition of primary end point (POI)	No resumption of menstrual activity and postmenopausal levels of FSH and E2 at 12 months after chemotherapy	Amenorrhea for the prior 6 months and postmenopausal levels of FSH at 24 months after chemotherapy	Amenorrhea between 12 and 24 months after random assignment with elevated FSH	No reappearance of two consecutive menstrual periods within 21 to 35 days 6 months after chemotherapy	No maintenance of menses and no resumption of menses at 24 months after chemotherapy
Planned sample size	280	416	250	62	124
Actual sample size	281	257	227	60	48
Start of accrual	October 2003	February 2004	August 2004	March 2005	July 2003
End of accrual	January 2008	May 2011	December 2009	December 2007	January 2007
ER status for eligibility	ER-positive and ER-negative	ER-negative only	ER-positive and ER-negative	ER-negative only	ER-positive and ER-negative
Upper age limit for eligibility	≤ 45 years	≤ 49 years	None	≤ 45 years	≤ 44 years
Stage for eligibility	I-III	I-III A	I-III B	I-III	I-III
Type of GnRH _a used in the experimental group	Triptorelin	Goserelin	Goserelin	Goserelin	Triptorelin
Mode of GnRH _a administration in the experimental group	3.75 mg IM started at least 1 week before chemotherapy and then every 4 weeks	3.6 mg SC started at least 1 week before chemotherapy and then every 4 weeks	3.6 mg SC implant started at least 1 week before chemotherapy and then every 4 weeks	3.6 mg SC started at least 2 weeks before chemotherapy and then every 4 weeks	3.75 mg IM started at least 1 week before chemotherapy and then every 4 weeks
NCI-CTCAE version	Version 2.0	Version 3.0	AE not recorded	Version 2.0	AE not recorded

Abbreviations: AC, doxorubicin, cyclophosphamide; AC/EC→D, AC or EC followed by docetaxel; AC→T, AC followed by a taxane; AE, adverse events; CAF, cyclophosphamide, doxorubicin, fluorouracil; CEF, cyclophosphamide, epirubicin, fluorouracil; CMF, cyclophosphamide, methotrexate, fluorouracil; E2, estradiol; EC, epirubicin, cyclophosphamide; E→CMF, epirubicin followed by CMF; EC→P, EC followed by paclitaxel; ED, epirubicin, docetaxel; ED→CMF, ED followed by CMF; EP→CMF, epirubicin, paclitaxel followed by CMF; ER, estrogen receptor; FAC, fluorouracil, doxorubicin, cyclophosphamide; FEC, fluorouracil, epirubicin, cyclophosphamide; FEC→D, FEC followed by docetaxel; FEC→GEM, FEC followed by gemcitabine; FEC→P, FEC followed by paclitaxel; FSH, follicle-stimulating hormone; GBG-37 ZORO, German Breast Group-37 ZOladex Rescue of Ovarian function; GnRH_a, gonadotropin-releasing hormone agonists; IM, intramuscularly; NCI-CTCAE, National Cancer Institute–Common Terminology Criteria for Adverse Events; NR, not reported; OPTION, Ovarian Protection Trial in Oestrogen Non-responsive Premenopausal Breast Cancer Patients Receiving Adjuvant or Neo-adjuvant Chemotherapy; POEMS, Prevention Of Early Menopause Study; POI, premature ovarian insufficiency; PROMISE-GIM6, PRevention Of Menopause Induced by chemotherapy: A Study in Early breast cancer patients—Gruppo Italiano Mammella 6; SC, subcutaneous; TAC, docetaxel, doxorubicin, cyclophosphamide.

eight trials were not available because of refusal to participate for three trials and no success in reaching the principal investigators of the other five trials despite several attempts to contact them.

Table 1 summarizes the main characteristics of the five included trials. As per study inclusion criteria, only patients with hormone receptor–negative breast cancer were enrolled in two trials.^{11,12}

A total of 873 patients were included, of whom 436 were randomly assigned to the GnRHa group and 437 to the control group (Data Supplement). Baseline patient and treatment characteristics were well balanced between the two groups (Table 2). Median age at diagnosis was 38 years (IQR, 34–42 years); 350 patients (40.1%) had estrogen receptor–positive disease.

Efficacy Results: Preservation of Ovarian Function and Fertility

Chemotherapy-induced POI was the primary end point in all trials; different definitions and time points for its evaluation were used (Table 1). POI data were available in 722 (82.7%) of 873 patients. In the GnRHa group, 51 (14.1%) of 363 patients developed POI, as compared with 111 (30.9%) of 359 in the control group (adjusted OR, 0.38; 95% CI, 0.26 to 0.57; $P < .001$). The meta-analysis approach showed no heterogeneity ($I^2 = 0\%$; $P = .726$; Fig 1A). The effect of GnRHa on reducing the risk of developing chemotherapy-induced POI was homogeneous among the different patient subgroups (Fig 1B). Multivariate analysis

showed that only treatment with GnRHa (adjusted OR, 0.38; 95% CI, 0.26 to 0.57; $P < .001$) and younger age at diagnosis (adjusted OR, 0.35; 95% CI, 0.24 to 0.52; $P < .001$) were significantly associated with a reduced risk of developing chemotherapy-induced POI (Data Supplement).

One-year amenorrhea data were available in 760 (87.1%) of 873 patients. In the GnRHa group, 142 (36.8%) of 386 patients developed 1-year amenorrhea as compared with 151 (40.4%) of 374 in the control group (adjusted OR, 0.92; 95% CI, 0.66 to 1.28; $P = .623$; Data Supplement).

Two-year amenorrhea data were available in 424 (48.6%) of 873 patients; this end point was not collected in the PROMISE-GIM6 study,^{9,10} and in patients who developed DFS and/or OS events between 1 and 2 years. In the GnRHa group, 39 (18.2%) of 214 patients developed 2-year amenorrhea, as compared with 63 (30.0%) of 210 in the control group (adjusted OR, 0.51; 95% CI, 0.31 to 0.85; $P = .009$; Data Supplement).

The three largest trials reported post-treatment pregnancies^{7,9–11}; preservation of fertility was a preplanned secondary end point in only one trial.¹¹ Information on post-treatment pregnancies was available in 726 (83.2%) of 873 patients. In the GnRHa group, 37 (10.3%) of 359 women had at least one post-treatment pregnancy, as did 20 (5.5%) of 367 in the control group (IRR, 1.83; 95% CI, 1.06 to 3.15; $P = .030$; Data Supplement). The meta-analysis approach showed no heterogeneity ($I^2 = 0\%$; $P = .852$; Fig 2). All pregnancies occurred in patients with ≤ 40 years of age at the time of diagnosis; 49 (86.0%) and eight (14.0%) of 57 were

Table 2. Baseline Patient and Tumor Characteristics and Treatments Administered by Study Group (N = 873)

Characteristic or Treatment	GnRHa Group (n = 436) No. (%)	Control Group (n = 437) No. (%)	P*
Age, years, median (IQR)	38 (34–42)	39 (35–42)	.258
Age distribution, years			.316
≤ 40	297 (68.1)	283 (64.8)	
≥ 41	139 (31.9)	154 (35.2)	
Tumor stage			.800
Stage I	76 (17.4)	78 (17.8)	
Stage II	134 (30.7)	130 (29.7)	
Stage III	62 (14.2)	54 (12.4)	
Missing	164 (37.6)	175 (40.0)	
Estrogen receptor status			.782
Positive	177 (40.6)	173 (39.6)	
Negative	257 (58.9)	262 (59.9)	
Missing	2 (0.5)	2 (0.5)	
Type of chemotherapy			.196
Anthracycline only	194 (44.5)	198 (45.3)	
Anthracycline plus taxane	227 (52.1)	210 (48.0)	
Non-anthracycline	6 (1.4)	13 (3.0)	
Missing	9 (2.1)	16 (3.7)	
Cumulative cyclophosphamide dose, mg/m ² , median (IQR)	4,000 (3,420 to 5,185)	3,960 (3,082 to 5,400)	.585
Duration of chemotherapy			.870
≤ 4 months	120 (27.5)	116 (26.5)	
> 4 months	208 (47.7)	194 (44.4)	
Missing	108 (24.8)	127 (29.1)	
Use of adjuvant endocrine therapy†			.578
No	8 (4.5)	5 (2.9)	
Yes	126 (71.2)	116 (67.0)	
Missing	43 (24.3)	52 (30.1)	

Abbreviations: GnRHa, gonadotropin-releasing hormone agonists; IQR, interquartile range.

*Calculated by excluding unknown data

†The percentages were calculated on the total number of patients with estrogen receptor–positive disease (177 in the GnRHa group and 173 in the control group).




Fig 1. Premature ovarian insufficiency (A) by trial, and (B) by patient subgroup. GBG-37 ZORO, German Breast Group-37 Zoladex Rescue of Ovarian function; GnRHa, gonadotropin-releasing hormone agonists; OPTION, Ovarian Protection Trial in Oestrogen Non-responsive Premenopausal Breast Cancer Patients Receiving Adjuvant or Neo-adjuvant Chemotherapy; OR, odds ratio; POEMS, Prevention Of Early Menopause Study; PROMISE-GIM6, PREvention Of Menopause Induced by chemotherapy: a Study in Early breast cancer patients—Gruppo Italiano Mammella 6; pts, patients.

observed in women with estrogen receptor–negative and estrogen receptor–positive disease, respectively (Data Supplement).

Safety Results: Toxicity and Survival Outcomes

GnRHa-related adverse events of any grade were recorded in three studies,⁹⁻¹² and two of them reported their severity by grade.⁹⁻¹¹

Concurrent administration of GnRHa and chemotherapy was associated with a significantly higher incidence of hot flashes and sweating, of grade 1 or 2 in the majority of the cases (Data Supplement). No significant difference was observed in the incidence of mood changes, vaginal dryness, and headache between the GnRHa and control groups.

Survival outcomes were collected in all included trials but one.¹³ Median follow-up time was 5.0 years (IQR, 3.0-6.3 years).




Fig 2. Post-treatment pregnancies by trial. GnRHa, gonadotropin-releasing hormone agonists; IRR, incidence rate ratio; OPTION, Ovarian Protection Trial in Oestrogen Non-responsive Premenopausal Breast Cancer Patients Receiving Adjuvant or Neo-adjuvant Chemotherapy; POEMS Prevention Of Early Menopause Study; PROMISE-GIM6, PRvention Of Menopause Induced by chemotherapy: a Study in Early breast cancer patients—Gruppo Italiano Mammella 6; pts, patients.

Among the 809 (92.7%) of 873 patients evaluable for DFS, 136 events (16.8%) were observed, 69 (17.2%) of 402 in the GnRHa group and 67 (16.5%) of 407 in the control group. Five-year DFS was 79.5% (95% CI, 74.7% to 83.5%) in the GnRHa group and 80.0% (95% CI, 75.2% to 83.9%) in the control group (adjusted HR, 1.01; 95% CI, 0.72 to 1.42; $P = .999$; Fig 3A). The meta-analysis approach showed low heterogeneity ($I^2 = 3.1\%$; $P = .377$; Data Supplement). Subgroup analysis according to estrogen receptor status showed no significant interaction ($P_{\text{interaction}} = .867$); the adjusted HRs were 1.17 (95% CI, 0.62 to 2.20) and 0.95 (95% CI, 0.64 to 1.42) in patients with estrogen receptor–positive (Fig 3B) and estrogen receptor–negative (Fig 3C) disease, respectively. The Data Supplement reports the multivariate analysis for DFS.

Among the 812 (93.0%) of 873 patients evaluable for OS, 77 events (9.5%) were observed, 33 (8.2%) of 404 in the GnRHa group and 44 (10.8%) of 408 in the control group. Five-year OS was 90.2% (95% CI, 86.4% to 92.9%) in the GnRHa group and 86.3% (95% CI, 82.0% to 89.7%) in the control group (adjusted HR, 0.67; 95% CI, 0.42 to 1.06; $P = .083$; Fig 4A). The meta-analysis approach showed that heterogeneity was rather high ($I^2 = 51.1\%$; $P = .105$; Data Supplement). Subgroup analysis according to estrogen receptor status showed no significant interaction ($P_{\text{interaction}} = .762$); the adjusted HRs were 0.79 (95% CI, 0.24 to 2.59) and 0.65 (95% CI, 0.39 to 1.07) in patients with estrogen receptor–positive (Fig 4B) and estrogen receptor–negative (Fig 4C) disease, respectively. The Data Supplement reports the multivariate analysis for OS.

DISCUSSION

This meta-analysis included individual patient–level data from five major trials that investigated the role of temporary ovarian suppression with GnRHa during chemotherapy as a strategy to preserve ovarian function and fertility in premenopausal women with early breast cancer. Concurrent administration of GnRHa and chemotherapy significantly reduced the risk of developing chemotherapy-induced POI and was associated with a higher

number of post-treatment pregnancies. No heterogeneity of treatment effect between subgroups was shown. Similar DFS and OS were observed between groups irrespective of the estrogen receptor status of the disease.

The avoidance of symptoms associated with chemotherapy-induced loss of gonadal function represents an important goal to be achieved in young survivors of breast cancer, even in patients without the desire to have a subsequent pregnancy.²² Our study supports the protective gonadal effect of GnRHa administration during chemotherapy, with a significant 16.8% absolute reduction in the incidence of chemotherapy-induced POI (adjusted OR, 0.38; 95% CI, 0.26 to 0.57; $P < .001$). The efficacy of this strategy was consistent across all subgroups analyzed, including in patients with estrogen receptor–positive disease, and independently of their age at the time of treatment. The different definition and time point of evaluation used in the trials highlights the current lack of standardized definition of chemotherapy-induced POI. Nevertheless, the availability of individual patient–level data allowed analysis of ovarian function recovery on the basis of more homogeneous definitions. Using 1-year amenorrhea, the reduced absolute 3.6% difference favoring the GnRHa was not statistically significant (adjusted OR, 0.92; 95% CI, 0.66 to 1.28; $P = .623$). On the contrary, our prior meta-analysis on the basis of abstracted data showed a significant reduction in the risk of 1-year amenorrhea (OR, 0.55; 95% CI, 0.41 to 0.73; $P < .001$) when considering the eight trials that reported this end point.⁶ The more limited number of trials included in the current analysis may explain this discrepancy. Nevertheless, the benefit of concurrent administration of GnRHa and chemotherapy became clearly evident at a longer time point, 2 years after the end of chemotherapy (11.8% absolute reduction; adjusted OR, 0.51; 95% CI, 0.31 to 0.85; $P = .009$). However, this end point could be evaluated in a more limited number of patients as compared with 1-year amenorrhea. This is likely attributable to longer-term evaluation not being originally planned for most of these studies and the difficulty faced by investigators in collecting information on end points like menstrual function. These findings also highlight that even if the majority of patients experience menstrual function recovery in the first




Fig 3. Disease-free survival (DFS) in (A) the whole study population, (B) patients with estrogen receptor–positive disease, and (C) patients with estrogen receptor–negative disease. GnRH α , gonadotropin-releasing hormone agonists.

12 months after chemotherapy, this can also occur beyond 1 year. Hence, an evaluation of chemotherapy-induced POI too close to the end of chemotherapy might not have captured the protective effect of GnRH α that became more evident at a longer time point. These findings also support the recent expert opinion–based suggestion to define menopausal status after chemotherapy not earlier than 2 years after the end of treatment.¹

Approximately 50% of young patients with breast cancer are concerned about the possible risk of infertility as a consequence of chemotherapy use and desire to have children after the end of treatment.²³ Nevertheless, these patients are, among survivors of cancer, those with the lowest chances of subsequent pregnancies.⁴ In our analysis, less than 10% of the patients had a post-treatment pregnancy (57 patients, 7.9%), in line with the available data in the literature of a pregnancy rate in survivors of breast cancer ranging between 4% and 7%.²⁴ Although the absolute numbers remain low and the trials were not designed to address pregnancy as primary end point, a statistically significant higher number of patients who underwent temporary ovarian suppression with GnRH α during

chemotherapy had a subsequent pregnancy as compared with those who received cytotoxic therapy alone (37 *v* 20; IRR, 1.83; 95% CI, 1.06 to 3.15; *P* = .030). This suggests the potential role of this strategy also as a fertility preservation procedure. Embryo and oocyte cryopreservation remain the first options to be proposed to women interested in fertility preservation.^{4,5} Nevertheless, being not mutually exclusive, temporary ovarian suppression with GnRH α during chemotherapy can also be used in this setting after cryopreservation procedures to increase the chances of a subsequent pregnancy as well as in patients who do not have access to assisted reproductive options.²⁵ Notably, for patients receiving temporary ovarian suppression with GnRH α during chemotherapy after prior controlled ovarian stimulation for embryo/oocyte cryopreservation, the timing for administering the long-acting GnRH α needs to be further explored, considering its potential use as trigger of final follicular maturation instead of chorionic gonadotropin or short-acting GnRH α . In addition, future research efforts should aim at clarifying the fertility outcomes of patients who undergo temporary ovarian suppression with GnRH α during




Fig 4. Overall survival (OS) in (A) the whole study population, (B) patients with estrogen receptor–positive disease, and (C) patients with estrogen receptor–negative disease. GnRH α , gonadotropin-releasing hormone agonists.

chemotherapy after cryopreservation procedures as compared with those of women who access only one of the two strategies.²⁵

The main adverse events associated with GnRH α administration are vasomotor symptoms and sexual problems.²⁶ Nevertheless, these side effects are mainly of grade 1 or 2 and are reversible at the time of treatment completion. Importantly, preservation of ovarian function with GnRH α use during chemotherapy may help avoid menopausal symptoms, including loss of bone density, in the long term; this is of crucial importance also in women not interested in fertility preservation.

In the past, two major safety concerns on the use of temporary ovarian suppression with GnRH α during chemotherapy were raised for women with estrogen receptor–positive disease: a potential antagonism with concurrent administration of antiestrogen therapy and cytotoxic systemic therapy, and the possible detrimental effect on prognosis of the lack of chemotherapy-induced POI.²⁷ These concerns have been recently dispelled by the results of the TEXT (Tamoxifen and Exemestane Trial) and SOFT (Suppression of Ovarian Function Trial) trials showing no survival

difference between patients who received GnRH α concurrently or sequentially to chemotherapy.²⁸ Our study confirms the safety of concurrent administration of GnRH α and chemotherapy in all patients with breast cancer, irrespective of the estrogen receptor status of their disease. Nevertheless, of note, the majority of patients with estrogen receptor–positive disease included in this analysis derive from the PROMISE-GIM6 trial.^{9,10} In this study, approximately 70% of the patients with estrogen receptor–positive disease who had resumed ovarian function after chemotherapy were treated with GnRH α as part of adjuvant endocrine therapy.¹⁰ This further supports that the safety of ovarian function preservation in patients with hormone receptor–positive disease should be considered in the context of subsequent ovarian function suppression as part of adjuvant endocrine therapy.²⁹

Some limitations of the current study should be acknowledged. It was possible to include individual patient–level data from only five major randomized trials, corresponding to 55.2% (873 of 1,581) of the eligible population for this study.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at jco.org.

AUTHOR CONTRIBUTIONS

Conception and design: Matteo Lambertini, Ann H. Partridge, Lucia Del Mastro

Provision of study materials or patients: Matteo Lambertini, Halle C.F. Moore, Robert C.F. Leonard, Sibylle Loibl, Pamela Munster, Richard A. Anderson, Susan Minton, Francesca Poggio, Kathy S. Albain, Douglas J.A. Adamson, Bernd Gerber, Gianfilippo Bertelli, Sabine Seiler, Amy Cripps, Ann H. Partridge, Lucia Del Mastro

Collection and assembly of data: Matteo Lambertini, Halle C.F. Moore, Robert C.F. Leonard, Sibylle Loibl, Pamela Munster, Marco Bruzzone, Luca Boni, Joseph M. Unger, Richard A. Anderson, Keyur Metha, Francesca Poggio, Kathy S. Albain, Douglas J.A. Adamson, Bernd Gerber, Gianfilippo Bertelli, Sabine Seiler, Marcello Ceppi, Ann H. Partridge, Lucia Del Mastro

Data analysis and interpretation: Matteo Lambertini, Marco Bruzzone, Marcello Ceppi, Ann H. Partridge, Lucia Del Mastro

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

Nevertheless, for all end points that could be evaluated also in our previous meta-analysis of abstracted data, similar results were observed, with the exception of 1-year amenorrhea.⁶ This may suggest that the inclusion of all studies would not have modified the overall findings. For the efficacy end points, major limitations are the lack of data on the extent of ovarian function preservation using more sensitive biomarkers like the anti-Müllerian hormone, the limited information on patients' wish to have a pregnancy, and on the number of women with more than one post-treatment pregnancy. Finally, a few baseline characteristics and end point data were missing in some trials and for some patients also in studies that had originally planned to collect them. Nevertheless, there is no evidence of imbalance for these missing data or a differential drop-out between the GnRHa and control groups; thus, it is unlikely that they may have influenced the comparisons between randomly assigned groups.

In conclusion, our systematic review and meta-analysis of individual patient-level data provides evidence for the efficacy and safety of temporary ovarian suppression with GnRHa during chemotherapy in premenopausal patients with early breast cancer. Given the findings of our study, this strategy should be considered as an available option to reduce the likelihood of chemotherapy-induced POI and potentially improve future fertility in premenopausal patients with early breast cancer undergoing (neo)-adjuvant chemotherapy.

REFERENCES

- Paluch-Shimon S, Pagni O, Partridge AH, et al: ESO-ESMO 3rd international consensus guidelines for breast cancer in young women (BCY3). *Breast* 35:203-217, 2017
- Lambertini M, Goldrat O, Clatot F, et al: Controversies about fertility and pregnancy issues in young breast cancer patients: Current state of the art. *Curr Opin Oncol* 29:243-252, 2017
- Howard-Anderson J, Ganz PA, Bower JE, et al: Quality of life, fertility concerns, and behavioral health outcomes in younger breast cancer survivors: A systematic review. *J Natl Cancer Inst* 104:386-405, 2012
- Peccatori FA, Azim HA Jr, Orecchia R, et al: Cancer, pregnancy and fertility: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 24:vi160-vi170, 2013 (suppl 6)
- Loren AW, Mangu PB, Beck LN, et al: Fertility preservation for patients with cancer: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol* 31:2500-2510, 2013
- Lambertini M, Ceppi M, Poggio F, et al: Ovarian suppression using luteinizing hormone-releasing hormone agonists during chemotherapy to preserve ovarian function and fertility of breast cancer patients: A meta-analysis of randomized studies. *Ann Oncol* 26:2408-2419, 2015
- Leonard RCF, Adamson DJA, Bertelli G, et al: GnRH agonist for protection against ovarian toxicity during chemotherapy for early breast cancer: The Anglo Celtic Group OPTION trial. *Ann Oncol* 28:1811-1816, 2017
- Moher D, Liberati A, Tetzlaff J, et al: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 339:b2535, 2009
- Del Mastro L, Boni L, Michelotti A, et al: Effect of the gonadotropin-releasing hormone analogue triptorelin on the occurrence of chemotherapy-induced early menopause in premenopausal women with breast cancer: A randomized trial. *JAMA* 306:269-276, 2011
- Lambertini M, Boni L, Michelotti A, et al: Ovarian suppression with triptorelin during adjuvant breast cancer chemotherapy and long-term ovarian function, pregnancies, and disease-free survival: A randomized clinical trial. *JAMA* 314:2632-2640, 2015
- Moore HCF, Unger JM, Phillips K-A, et al: Goserelin for ovarian protection during breast-cancer adjuvant chemotherapy. *N Engl J Med* 372:923-932, 2015
- Gerber B, von Minckwitz G, Stehle H, et al: Effect of luteinizing hormone-releasing hormone agonist on ovarian function after modern adjuvant breast cancer chemotherapy: The GBG 37 ZORO study. *J Clin Oncol* 29:2334-2341, 2011
- Munster PN, Moore AP, Ismail-Khan R, et al: Randomized trial using gonadotropin-releasing hormone agonist triptorelin for the preservation of ovarian function during (neo)adjuvant chemotherapy for breast cancer. *J Clin Oncol* 30:533-538, 2012
- Badawy A, Elnashar A, El-Ashry M, et al: Gonadotropin-releasing hormone agonists for prevention of chemotherapy-induced ovarian damage: Prospective randomized study. *Fertil Steril* 91:694-697, 2009
- Sverrisdottir A, Nystedt M, Johansson H, et al: Adjuvant goserelin and ovarian preservation in chemotherapy treated patients with early breast cancer: Results from a randomized trial. *Breast Cancer Res Treat* 117:561-567, 2009
- Elgindy EA, El-Haieg DO, Khorshid OM, et al: Gonadotrophin suppression to prevent chemotherapy-induced ovarian damage: A randomized controlled trial. *Obstet Gynecol* 121:78-86, 2013
- Song G, Gao H, Yuan Z: Effect of leuprolide acetate on ovarian function after cyclophosphamide-doxorubicin-based chemotherapy in premenopausal patients with breast cancer: Results from a phase II randomized trial. *Med Oncol* 30:667, 2013
- Karimi-Zarchi M, Forat-Yazdi M, Vafaenasab MR, et al: Evaluation of the effect of GnRH agonist on menstrual reverse in breast cancer cases treated with cyclophosphamide. *Eur J Gynaecol Oncol* 35:59-61, 2014
- Sun J, Ren Y, Li W: Effect of zoladex administered before chemotherapy on menstruation of patients with breast cancer. *China Disability Medicine* 19:15-16, 2011
- Li JW, Liu GY, Yu KD, et al: Effect of using LHRH analog during chemotherapy (CT) on premature ovarian failure and prognosis in premenopausal patients with early-stage, hormone receptor-positive breast cancer: The primary analysis of a randomized controlled phase III trial. *Cancer Res* 75: P1-12-02, 2015 (abstr)
- Li M, Huang H, Liang Y, et al: Effect of zoladex administered before chemotherapy on menstruation of patients with breast cancer. *Chinese Journal of Clinical Oncology* 35:905-907, 2008
- Rosenberg SM, Tamimi RM, Gelber S, et al: Treatment-related amenorrhea and sexual functioning in young breast cancer survivors. *Cancer* 120:2264-2271, 2014
- Ruddy KJ, Gelber SI, Tamimi RM, et al: Prospective study of fertility concerns and preservation strategies in young women with breast cancer. *J Clin Oncol* 32:1151-1156, 2014
- Azim HA Jr, Kroman N, Paesmans M, et al: Prognostic impact of pregnancy after breast cancer according to estrogen receptor status: A multicenter retrospective study. *J Clin Oncol* 31:73-79, 2013
- Lambertini M, Cinquini M, Moschetti I, et al: Temporary ovarian suppression during chemotherapy

to preserve ovarian function and fertility in breast cancer patients: A GRADE approach for evidence evaluation and recommendations by the Italian Association of Medical Oncology. *Eur J Cancer* 71:25-33, 2017

26. Ribi K, Luo W, Bernhard J, et al: Adjuvant tamoxifen plus ovarian function suppression versus tamoxifen alone in premenopausal women with early breast cancer: Patient-reported outcomes in the

suppression of ovarian function trial. *J Clin Oncol* 34:1601-1610, 2016

27. Rugo HS, Rosen MP: Reducing the long-term effects of chemotherapy in young women with early-stage breast cancer. *JAMA* 306:312-314, 2011

28. Regan MM, Walley BA, Francis PA, et al: Concurrent and sequential initiation of ovarian function suppression with chemotherapy in premenopausal

women with endocrine-responsive early breast cancer: An exploratory analysis of TEXT and SOFT. *Ann Oncol* 28:2225-2232, 2017

29. Burstein HJ, Lacchetti C, Anderson H, et al: Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: American Society of Clinical Oncology clinical practice guideline update on ovarian suppression. *J Clin Oncol* 34:1689-1701, 2016

Affiliations

Matteo Lambertini, Institut Jules Bordet, Université Libre de Bruxelles, Brussels, Belgium; **Halle C.F. Moore**, Cleveland Clinic Taussig Cancer Institute, Cleveland, OH; **Robert C.F. Leonard**, Imperial College, London; **Richard A. Anderson**, University of Edinburgh, Edinburgh; **Douglas J.A. Adamson**, Ninewells Hospital, Dundee; **Gianfilippo Bertelli**, Sussex Cancer Centre, Brighton, United Kingdom; **Sibylle Loibl**, **Keyur Mehta**, and **Sabine Seiler**, German Breast Group, Neu-Isenburg; **Bernd Gerber**, University Hospital Rostock, Rostock, Germany; **Pamela Munster**, University of California, San Francisco, San Francisco, CA; **Marco Bruzzone**, **Francesca Poggio**, **Marcello Ceppi**, and **Lucia Del Mastro**, Ospedale Policlinico San Martino; **Francesca Poggio** and **Lucia Del Mastro**, University of Genova, Genova; **Luca Boni**, Azienda Ospedaliero Universitaria Careggi, Florence, Italy; **Joseph M. Unger**, SWOG Statistical Center and Fred Hutchinson Cancer Research Center, Seattle, WA; **Susan Minton**, Moffitt Cancer Center, Tampa, FL; **Kathy S. Albain**, Loyola University Chicago Stritch School of Medicine, Maywood, IL; **Amy Cripps**, Nexgen Oncology, Dallas, TX; and **Ann H. Partridge**, Dana-Farber Cancer Institute, Boston, MA.

Support

Supported in part by the Italian Association for Cancer Research (AIRC) Grant No. 2013:14272.

The financial sponsor of the study (AIRC) had no role in study design, data collection, or analysis, interpretation, or writing of the report, and it had no access to the individual patient-level data.

Prior Presentation

Presented at the 2017 San Antonio Breast Cancer Symposium, San Antonio, TX, December 7, 2017.



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Gonadotropin-Releasing Hormone Agonists During Chemotherapy for Preservation of Ovarian Function and Fertility in Premenopausal Patients With Early Breast Cancer: A Systematic Review and Meta-Analysis of Individual Patient–Level Data

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/site/ifc.

Matteo Lambertini

Consulting or Advisory Role: Teva

Travel, Accommodations, Expenses: Astellas Pharma

Halle C.F. Moore

Research Funding: Puma Biotechnology (Inst), AbbVie (Inst)

Robert C.F. Leonard

No relationship to disclose

Sibylle Loibl

Research Funding: Pfizer, Roche, Celgene, Amgen, Novartis (Inst)

Pamela Munster

Honoraria: Sanofi, Amgen (I), AstraZeneca, EMD Serono

Consulting or Advisory Role: OncoSec

Research Funding: Merck (Inst), Pfizer (Inst), Novartis (Inst), GlaxoSmithKline (Inst), OncoMed (Inst), Celgene (Inst), Andes Biotechnologies (Inst), Incyte (Inst), Ignyta (Inst), CBT Pharmaceuticals (Inst), Merrimack (Inst), Genentech (Inst), OncoSec (Inst), Bristol-Myers Squibb (Inst), Plexxikon (Inst), Piramal Life Science (Inst), Andes Biotechnologies (Inst), Immune Design (Inst), Biomarin (Inst)

Marco Bruzzone

No relationship to disclose

Luca Boni

No relationship to disclose

Joseph M. Unger

No relationship to disclose

Richard A. Anderson

Consulting or Advisory Role: Roche Diagnostics, HRA Pharma, NeRRe Therapeutics

Speakers' Bureau: IBSA, Merck

Research Funding: Ferring Pharmaceuticals

Keyur Metha

No relationship to disclose

Susan Minton

No relationship to disclose

Francesca Poggio

No relationship to disclose

Kathy S. Albain

Consulting or Advisory Role: Novartis, Pfizer, Myriad Genetics, Genentech

Other Relationship: Puma Biotechnology

Douglas J.A. Adamson

Stock or Other Ownership: GlaxoSmithKline (I)

Research Funding: Roche (Inst), Aventis Pharma (Inst), Novartis (Inst), Amgen (Inst), Pfizer (Inst), Bayer (Inst), Sanofi (Inst), Boehringer Ingelheim (Inst), Immodulon Therapeutics (Inst), Schering-Plough (Inst), Biocompatibles (Inst), AstraZeneca (Inst)

Bernd Gerber

Travel, Accommodations, Expenses: AstraZeneca, Roche

Amy Cripps

Employment: NexGen Oncology

Travel, Accommodations, Expenses: Foundation Medicine

Gianfilippo Bertelli

Consulting or Advisory Role: Roche, Novartis Pharmaceuticals UK, Genomic Health

Travel, Accommodations, Expenses: Roche

Sabine Seiler

Consulting or Advisory Role: Hexal, Roche, Amgen, Novartis

Travel, Accommodations, Expenses: Hexal, Roche, Novartis, Amgen

Marcello Ceppi

No relationship to disclose

Ann H. Partridge

No relationship to disclose

Lucia Del Mastro

Honoraria: Roche, Novartis, Takeda Pharmaceuticals, Ipsen, Eisai, Pfizer

Consulting or Advisory Role: Eli Lilly

Acknowledgment

Matteo Lambertini acknowledges the support from the European Society for Medical Oncology (ESMO) for a Translational Research Fellowship at Institut Jules Bordet in Brussels (Belgium); he received a San Antonio Breast Cancer Symposium Clinical Scholar Award for presenting the results of this systematic review and meta-analysis at the 2017 San Antonio Breast Cancer Symposium. The authors thank all the patients and the investigators from the Gruppo Italiano Mammella (GIM) study group, the SWOG, the Anglo Celtic Group, the German Breast Group (GBG), and the Moffitt-led study who participated in the five included trials.